









- [6] 毕福强, 樊学忠, 许诚, 等. 不敏感四唑非金属含能离子化合物的研究进展[J]. 含能材料, 2012, 20(6): 805–811.  
BI Fu-qiang, FAN Xue-zhong, XU Cheng, et al. Review on insensitive non-metallic energetic ionic compounds of tetrazolate anions[J]. *Chinese Journal of Energetic Materials(Hanneng Cailiao)*, 2012, 20(6): 805–811.
- [7] Zhao X X, Li S H, Wang Y, et al. Design and synthesis energetic materials towards high density and positive oxygen balance by *N*-dinitromethyl functionalization of nitroazoles[J]. *Journal of Materials Chemistry A*, 2016, 4(15): 5495–5504.
- [8] Donald H, Brooke J, David T, et al. Lattice potential energy estimation for complex ionic salts from density measurements[J]. *Inorganic Chemistry*, 2002, 41(9): 2364–2367.
- [9] Dorofeeva O V, Ryzhova O N, Suntsova M A. Accurate prediction of enthalpies of formation of organic azides by combining G4 theory calculations with an isodesmic reaction scheme[J]. *The Journal of Physical Chemistry A*, 2013, 117(31): 6835–6845.
- [10] Head M, Pople J A. MP2 energy evaluation by direct methods[J]. *Chemical Physics Letters*, 1988, 153(6): 503–506.
- [11] Martin J M. Ab initio total atomization energies of small molecules-towards the basis set limit[J]. *Chemical Physics Letters*, 1996, 259(5–6): 669–678.
- [12] Zhang X L, Gong X D. Theoretical studies on the energetic salts of substituted 3,3'-amino-*N,N'*-azo-1,2,4-triazoles; the role of functional groups[J]. *Journal of Chemical and Engineering Data*, 2015, 60(10): 2869–2878.
- [13] Golovina N I, Tikkov A N, Raevskii A V, et al. Kinetics and mechanism of phase transition in the crystals of 2,4,6-trinitrotoluene and benzotrifuroxane[J]. *Journal of Solid State Chemistry*, 1994, 113(2): 229–238.

## Synthesis and Properties of 5-Methyl-4-nitro-1*H*-pyrazol-3-(2*H*)-one and its Energetic Ion Compounds

DENG Mu-cong, WANG Yi, ZHANG Wen-quan, ZHANG Qing-hua

(Institute of Chemical Materials, CAEP, Mianyang 621999, China)

**Abstract:** A new energetic compound 5-methyl-4-nitro-1*H*-pyrazol-3-(2*H*)-one (MNPO) was synthesized with total yield of 68%, using ethyl 3-oxobutanoate as raw material. Corresponding ion salts were prepared via metathesis reaction and neutralization reaction by the reactions of MNPO with a series of high nitrogen cation. Their structures were characterized by single-crystal X-ray diffraction, Fourier transform infrared(FT-IR) spectroscopy, nuclear magnetic resonance(<sup>1</sup>H NMR, <sup>13</sup>C NMR) spectrometry, elemental analyses and other means. Their thermal decomposition temperatures were measured by thermogravimetry(TG) and differential scanning calorimetry(DSC). Their detonation performances were calculated with Explo5 v6.02 software. Results show that the crystal of MNPO belongs to orthorhombic system, and its space group is *Pbca* with crystal parameters of  $a=71495(18)$  nm,  $b=1.1639(3)$  nm,  $c=1.3834(3)$  nm,  $V=1.1512(5)$  nm<sup>3</sup> and  $Z=8$ . For energetic ion salts of MNPO with densities of 1.62–1.74 g·cm<sup>-3</sup>, their decomposition temperatures are in the range of 181–272 °C, theoretical detonation velocities are greater than 7000 m·s<sup>-1</sup> and detonation pressures are greater than 15 GPa. The measured impact and friction sensitivities are low, in which, the impact sensitivity of ammonium salt of MNPO is 28 J and the friction sensitivity 240 N.

**Key words:** synthesis; 5-methyl-4-nitro-1*H*-pyrazol-3-(2*H*)-one(MNPO); energetic ion compounds; single crystal structure

**CLC number:** TJ55; O62

**Document code:** A

**DOI:** 10.11943/j.issn.1006-9941.2017.08.005